



## King's Research Portal

DOI:

[10.1016/j.archoralbio.2018.05.001](https://doi.org/10.1016/j.archoralbio.2018.05.001)

*Document Version*

Peer reviewed version

[Link to publication record in King's Research Portal](#)

*Citation for published version (APA):*

Proctor, G. (2018). Diurnal rhythm and salivary electrolyte. *Archives of Oral Biology*.  
<https://doi.org/10.1016/j.archoralbio.2018.05.001>

### **Citing this paper**

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

### **General rights**

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

### **Take down policy**

If you believe that this document breaches copyright please contact [librarypure@kcl.ac.uk](mailto:librarypure@kcl.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.

## Accepted Manuscript

Title: Diurnal rhythm and salivary electrolyte

Author: Gordon Proctor

PII: S0003-9969(18)30141-9

DOI: <https://doi.org/10.1016/j.archoralbio.2018.05.001>

Reference: AOB 4160

To appear in: *Archives of Oral Biology*

Received date: 26-3-2018



Please cite this article as: Proctor G, Diurnal rhythm and salivary electrolyte, *Archives of Oral Biology* (2018), <https://doi.org/10.1016/j.archoralbio.2018.05.001>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title is Diurnal rhythm and salivary electrolyte.

Auhtor is Gordon Proctor.

In the October, 2017 issue of Archives of Oral Biology a paper from Bel'skaya et al. was published describing an investigation into the influence of a diurnal rhythm on salivary mineral composition (Bel'skaya, Kosenok, & Sarf, 2017). Saliva samples were collected at 3 hr intervals over a 24 hr period and pH, calcium, phosphorus, sodium and potassium concentrations were assayed. Following publication, the Editors received a letter from Professor Colin Dawes, who has published seminal research papers on the influence of diurnal rhythms on saliva mineral composition, expressing a number of concerns regarding the paper.

In the present editorial we have highlighted the major concerns raised by Professor Dawes and in the interests of balance, have given the authors an opportunity to respond. The purpose of this editorial is two-fold. Firstly, it clarifies some aspects of the published paper and enables the reader to make a judgement based on the concerns raised and the answers given. Secondly, it highlights some aspects of salivary research that require the attention of those intending to undertake such studies.

Major concerns expressed by Professor Dawes:

1. The readers are not told specifically whether the saliva was unstimulated or stimulated, although it was probably unstimulated. The timing of meals in relation to the saliva collection times is also not given. This is important, since the composition of saliva collected within one hour of a meal may be changed (Dawes & Chebib, 1972).
2. The saliva samples were centrifuged prior to assay but as we are told only the revolutions per minute of the centrifuge and not its radius, it is not possible to calculate the centrifugal force applied in gravitational units.
3. All the assays, including measurement of the pH, were apparently done using a capillary electrophoresis system, with quantification of different components at different wavelengths. The description of the apparatus is incomprehensible, at least to this reader. Accurate measurement of pH requires that saliva not be exposed to the atmosphere, which causes a loss of CO<sub>2</sub> and a rise in pH. As no such precautions were mentioned, the pH values are suspect.
4. Prior to assay of various inorganic components of the saliva, salivary protein was said to be "deposited" (presumably precipitated) with 10% trichloroacetic acid. However, although this reagent precipitates most plasma proteins, it does not effectively precipitate most salivary proteins (Dawes, 1965).
5. The study began at 3 a.m. and saliva collections continued every three hours until midnight of the same day. However, in the four Figures, the midnight values have been duplicated at midnight of the previous day.
6. On the last two lines of page 287, it states that "higher salivary flow rates were associated with lower levels of sodium ions". This is the opposite of what would be expected, as the sodium concentration in saliva normally increases with flow rate.
7. The salivary potassium concentrations average about 9.5 mmol/L, which is about half the value expected from the literature.
8. The mean values at different time points for men alone, women alone and men + women in Figures 1 – 4, are all connected by beautifully-shaped curves which pass through every point. However, the derivation of these curves is not described.

9. On page 289, line 6 of column 1, it is written “The mineralizing function of saliva is carried out due to its supersaturation with  $\text{Ca}^{2+}$  and  $\text{HPO}_4^{2-}$  ions.” However, saliva can not be saturated with individual ions, only with respect to a mineral.
10. On page 290, line 5 of column 1, it is written “This finding can be explained through higher solubility of calcium phosphates at lower acidity of the medium.” This is nonsense, as the solubility of all calcium phosphates increases as the pH falls and decreases as it rises.

Responses from authors to comments:

1. Study plan included a collection of non-stimulated saliva not earlier than 1.5 hours after a meal.
2. Centrifugation was carried out at the acceleration (Relative Centrifugal Force) of 8610 g.
3. The capillary electrophoresis technique is widely described in the literature (Demkowska, Polkowska, & Namieśnik, 2008; Guo et al., 2016; Mori et al., 2016; Mori, Yamamoto, Kaseda, Yamada S, & Itabashi, 2012); this article provides an in-depth description of conditions for analysis and preparation of the samples (Bel'skaya, 2017). With regard to the pH measurement, saliva was collected into test tubes with tightly screwing caps, and pH was measured immediately after the opening of the tubes. Saliva samples were stored at a temperature of 2-8°C; all analyses were performed within 24 hours after collection.
4. Deposition of proteins with 10% trichloroacetic acid was carried out in order to prevent contamination of the capillary during the recording of electropherograms of the samples. Since protein composition of saliva was not analyzed, it could not affect the results and their interpretation.
5. There was no duplication of values, these data are the same and are shown in the graph.  
The experiment was repeated for 3 days, the values obtained at the same time are almost the same. The error does not exceed 10%.
6. Secretion of saliva is a two-step process: the first stage involves acini, the second – salivary ducts. Acini secrete primary secret that contains ptyalin and/or mucus in a solution with the concentration of ions that is not much different from the concentration of ions in extracellular fluid. As the primary secretion progresses through channels, sodium ions are actively reabsorbed by all salivary ducts and potassium ions are actively secreted in exchange for the sodium, so the concentration of sodium ions in saliva significantly reduces, while the concentration of potassium ions increases. In this article, we provided experimentally obtained fact that the sodium concentration and the salivation rate change in opposite directions. We did not touch upon this subject in our work, but a reduction of sodium level and an increase of potassium concentration may not only depend on the salivation rate (Jirakulsomchok & Schneyer, 1987). We hope to investigate this question more in our further studies.
7. In the literature, the concentration of potassium in saliva varies in a wide range. It may be due to the different conditions of collection and storage, as well as the modes of centrifugation and separation of a sediment that contains intracellular potassium from damaged cells giving the excessive concentrations.
8. Curves were plotted using interpolation method with Bezier curves (R Studio 3.3.0).
9. Basing on the theory of micellar structure of saliva  $\{[m(\text{Ca}_3(\text{PO}_4)_2]_n\text{HPO}_4^{2-}(n-x)\text{Ca}^{2+}\}^{2x-x\text{Ca}^{2+}}$ , the saliva is over-saturated with  $\text{Ca}^{2+}$  and  $\text{HPO}_4^{2-}$  in relation to the calcium phosphate.

10. The solubility of calcium phosphates decreases with increasing pH of the medium. This fact is confirmed by thermodynamic calculations, what is written on page 290, row 5, column 1. Here the solubility of calcium phosphates increases along with decreasing pH of the medium. We wanted to reflect this idea in our article. We want to say that this finding can be explained by higher solubility of calcium phosphates at lower pH of the medium.

## References

- Belskaya, L.V. (2017). Application of capillary electrophoresis for determination of the mineral composition of human saliva. *Bulletin of Science and Practice* 2: 132–140.
- Bel'skaya LV, Kosenok VK, Sarf EA. (2017) Chronophysiological features of the normal mineral composition of human saliva. *Archives of Oral Biology* 82: 286-292.
- Dawes C. (1965) Some characteristics of parotid and submandibular salivary proteins. *Archives of Oral Biology* 10: 269-78
- Dawes C, Chebib FS. (1972) The influence of previous stimulation and the day of the week on the concentrations of protein and the main electrolytes in human parotid saliva. *Archives of Oral Biology* 17: 1289-301.
- Demkowska I, Polkowska Z, Namieśnik J. (2008) Application of ion chromatography for the determination of inorganic ions, especially thiocyanates in human saliva samples as biomarkers of environmental tobacco smoke exposure. *Journal of Chromatography B*. 875: 419-26
- Guo L, Wang Y, Zheng Y, Huang Z, Cheng Y, Ye J, Chu Q, Huang D. (2016) Study on the potential application of salivary inorganic anions in clinical diagnosis by capillary electrophoresis coupled with contactless conductivity detection. *Journal of Chromatography B*. 1014:70-4.
- Jirakulsomchok, D., & Schneyer, C. (1987). Effects of  $\alpha$ ,  $\beta$ 1, and  $\beta$ 2 adrenergic antagonists on the Na and K concentrations of sympathetic-nerve stimulated rat saliva. *Journal of the Autonomic Nervous System*, 20, 65–71.
- Mori M, Ishikawara F, Tomoda T, Yamada S, Okamoto M, Itabashi H, Seki Y, Matsumoto R, Shoho Y, Martha L, Sumino H, Murakami M. (2016) Use of capillary electrophoresis with dual-opposite end injection for simultaneous analysis of small ions in saliva samples from wrestlers undergoing a weight training program. *Journal of Chromatography B* 1012-1013: 178-85
- Mori M, Yamamoto T, Kaseda M, Yamada S, Itabashi H. (2012) Capillary ion electrophoresis of inorganic anions and uric acid in human saliva using a polyvinyl alcohol coated capillary column and hexamethonium chloride as additive of background electrolyte. *Journal of Chromatography B* 887-888: 1-7.